

The carbon cycle

in the C-IFS model for atmospheric composition and weather prediction

Anna Agusti-Panareda

**Sebastien Massart, Mark Parrington,
Miha Ratzinger, Luke Jones, Michail Diamantakis
Gianpaolo Balsamo, Souhail Boussetta
Emanuel Dutra, Joaquin Munoz-Sabater,
Alessio Bozzo, Robin Hogan, Richard Forbes
(ECMWF)**

**Frederic Chevallier, Phillippe Peylin,
Natasha MacBean, Fabienne Maignan (LSCE)**

Anna.Agusti-Panareda@ecmwf.int



Funded by the European Union

Implemented by  ECMWF


The carbon cycle

Interaction between all the Earth system components

- Carbon reservoirs and their interactions with the atmosphere (focusing on CO₂ primarily).
- Can carbon cycle – climate feedbacks improve atmospheric predictive skill?

Vegetation, radiative transfer, atmospheric chemistry


- Atmospheric CO₂ and CH₄ analysis and forecast (Copernicus Service)



Funded by the European Union

The 'spheres' of influence on the climate system.
Source from [Institute for Computational Earth System Science\(ICESSE\)](http://www.icesse.org)

The atmospheric reservoir in the fast carbon cycle (annual time-scale)



Movement of carbon between land, atmosphere, and oceans:

Yellow numbers are natural (balanced fluxes)

Red are human contributions (perturbing balance)



[Units: in Gigatons of carbon per year]

White numbers: stored carbon [Gigatons of carbon].

Source: <http://earthobservatory.nasa.gov/Features/CarbonCycle/>
(Diagram adapted from U.S. DOE, Biological and Environmental Research Information System.)


The atmospheric reservoir:

surface observations



CO₂ growth rate in the atmospheric reservoir

The atmospheric concentration growth rate [Gt CO₂/year]



In 2015 CO₂ increased by 3 ppm
~ 23 GtCO₂/year:


(droughts associated and fires during el Nino episodes)

15 GtCO₂/year ~ 2 ppm/year on average for last 10 years


In 1997-1998 el Nino
CO₂ increased by 2.8 ppm

Global carbon budget

Data: CDIAC/NOAA-ESRL/GCP/Joos et al 2013/Khatiwala et al 2013





ANTHROPOGENIC FLUXES




EDGAR v4.2 inventory of anthropogenic emissions (excluding land-use change)

Source: EDGAR database



Source: [Global Carbon Budget 2015](#); CDIAC

CO₂ emissions: land-use change



CO₂ emissions: land-use change by burning biomass




GFAS daily fire product available 1 day behind real time


CAMS GFAS Daily Fire Products Sunday 04 September 2016

Average of Observed Fire Radiative Power Areal Density [mW/m²]

max value = 0.71 W/m²



Sept-Oct 2015 daily mean CO₂ emissions



GFAS CO₂ emissions over Indonesia (Sep-Oct 2015):


Fires contribute to el Nino signal in the atmospheric CO₂ growth rate

The ocean reservoir in the carbon cycle

Solubility pump
(inorganic carbon)

Ocean circulation
(long timescales)

Biological pump
(organic carbon)



The CO₂ ocean-atmosphere fluxes

Climatology of monthly mean ocean fluxes from Takahashi et al. (2009) used in C-IFS

Observations of pCO₂ at the surface of the ocean and in the atmosphere with transfer coefficients based on turbulent exchange.

Regions of sources and sinks associated with **upwelling** and **downwelling** regions




Fig. 13. Climatological mean annual sea-air CO₂ flux (g-C m⁻² yr⁻¹) for the reference year 2000 (non-El Niño conditions). The map is based on 3.0 million surface water pCO₂ measurements obtained since 1970. Wind speed data from the 1979–2005 NCEP-DOE AMIP-II Reanalysis (R-2) and the gas transfer coefficient with a scaling factor of 0.26 (Eq. (8)) are used. This yields a net global air-to-sea flux of 1.42 Pg-Cy⁻¹.

Takahashi et al. (2009)

The terrestrial CO₂ fluxes

- Strong link with water and energy fluxes





Figure 3. Mean annual (1982–2008) (a) GPP, (b) LE, (c) TER, and (d) H derived from global empirical upscaling of FLUXNET data.

Terrestrial carbon flux : Exchange between the biosphere and the atmosphere

Atmospheric CO₂ sink (Gross Primary Production):

Photosynthesis
(plants)




Atmospheric CO₂ source (Ecosystem Respiration):

Respiration
(plants,
animals)




+ decomposition of organic carbon in soil by microbes



Modelling CO₂ uptake by plants (GPP) in C-IFS

BOUSSETTA ET AL.: LAND CO₂ WITHIN THE ECMWF SYSTEM



$$r_s = \frac{1}{g_s}$$


$$g_s = \frac{A_n}{(C_s - C_i)}$$

Environmental factors:

- Temperature
- PAR (solar radiation)
- Soil moisture
- Atm. wv deficit
- Atm. CO₂

Biological factors:

- Mesophyll conductance



+ Soil moisture stress function


$$C_s - C_i = f(D_s, r_m)$$

$$C_s = 340 \text{ ppm}$$

CTESSEL parameterisation based on ISBA-Ags
 Jacobs (1994), Calvet et al., 1998,2000, Lafont et al. 2012, Boussetta et al. (2013)

Modelling CO₂ uptake by plants (GPP) in C-IFS


LEAF STOMATA



CANOPY



MODEL FLUXES




Upscaling to canopy with **LAI** climatology from MODIS


Upscaling to model grid point with **vegetation dominant type/cover**

Modelling soil respiration

$$R_{soil} = R_0 Q_{10}^{(0.1(T_{soil}-25))} f_{sm}$$



$$R_{soil} = R_0 e^{-\alpha \cdot Z_{snow}} Q_{10}^{(0.1(T_{soil}-25))} f_{sm}$$




Environmental factors:

- Temperature
- Soil moisture
- Snow depth


Biological factors:

- Organic carbon in soil and microbial activity (R0 parameter)



Including a snow attenuation effect on the soil CO2 emission


Q10 dependance on Temperature regime



Including a temperature dependancy on the Q10 parameter (McGuire et al., 1992)

Evaluation of CO₂ ecosystem fluxes from CTESSEL in IFS

Example of NEE (micro moles /m²/s) predicted over the site Fi-Hyy (FINLAND) by **CTESSEL (black line)** and **CASA-GFED3 (green-line)** compared to **FLUXNET observations**



Scheme	NEE rmse	NEE bias	NEE corr
CTESSEL	3.736	-1.656	0.536
CASA	1.872	0.739	0.297

Modelling atmospheric CO₂ in C-IFS

Synoptic variability of NEE is important for the CO₂ synoptic variability in the BL

In the warm sectors of low pressure systems:


synergy between advection and CO₂ ecosystem fluxes:

cloudy
warm

reduction of CO₂ uptake
increase in respiration

More CO₂

Enhanced atmospheric CO₂ anomaly



Modelling atmospheric CO₂ in C-IFS

CO₂ surface fluxes & column-averaged dry-air mole fraction of CO₂ [ppm]

Transport

IFS model

20150501 00 UTC

Fluxes

Vegetation (CTESSEL model)

Source ○

Sink ○



Fires (GFAS) ▲

Ocean (Takahashi et al 2009)

Source ○


Sink ○

Anthropogenic
(EDGAR v4.2) ●



Symbol size reflects the relative flux intensity
(Note that fires have been re-scaled by a factor of 10)



GOSAT analysis (28 November 2014 – 14 December 2014)



**Analysis departure (o-a)
In ppm for GOSAT data**



No or few GOSAT data to
constrain the analysis in
these regions

Massart et al. ACP 2015



Correcting atmospheric CO₂ biases with Biogenic Flux Adjustment Scheme (BFAS)

ARCHIVED DATA




EPS FORECAST/HINDCAST

BFAS

Compares budgets

Re-scaling maps for biogenic fluxes from CTESSEL



CAMS CO₂ modelling (IFS)

CO₂ SURFACE FLUXES

CTESSEL model

Prescribed

Anthropogenic emissions


Ocean sources/sinks

Fire emissions

Adjusted Net Ecosystem Fluxes




TRANSPORT

MACC column-averaged dry-air mole fraction of CO₂ (ppm)
September 2013



Improved atmospheric CO₂ forecast

Biogenic Flux Adjustment Scheme: Improving the total column CO₂



Total column mean TCCON Observations
Atmospheric CO₂ simulations with optimized fluxes
climatology of optimized fluxes
Modelled NEE
Modelled NEE + BFAS

Biogenic Flux Adjustment Scheme: Improving CO₂ synoptic variability



March 2010



**NOAA/ESRL tall tower
Observations**

*Atmospheric CO₂ simulations with
optimized fluxes*




climatology of optimized fluxes

Modelled NEE

Modelled NEE + BFAS




CO₂ Ecosystem Flux Adjustment factors: what can we learn to improve the model?



Month

Crops



Vegetation code	Vegetation type
1	Crops, mixed farming
2	Short grass
7	Tall grass
9	Tundra
10	Irrigated crops
11	Semidesert
13	Bogs and marshes
16	Evergreen shrubs
17	Deciduous shrubs
3	Evergreen needle leaf trees
4	Deciduous needle leaf Trees
5	Deciduous broadleaf trees
6	Evergreen broadleaf trees
18	Mixed forest/woodland
19	Interrupted forest
21	Tropical savanna (new type)
-	Remaining land points without veg



- Re-tune the reference respiration for crops
- Distinction between C3 and C4 crops necessary
- Revision of vegetation types: A new subtype of interrupted forest for BFAS (tropical savanna)





Feedbacks of carbon cycle to NWP:

- Improvement in representation of vegetation:
photosynthesis, phenology, albedo**

Jarvis Vs photosynthesis-based evapotranspiration (offline run)



Surface sensible heat flux (W/m^2) compared with flux-tower observations over Fr-LBr for HTESSEL (left panel) and CCTESSEL (right panel)




Surface latent heat flux (W/m^2) compared with flux-tower observations over Fr-LBr for HTESSEL (left panel) and CCTESSEL (right panel).

- **CCTESSEL improves the LE/H simulations (Photosynthesis-based vs Jarvis approach).**

LE/H: When “good” is not enough? (Interaction with the atmosphere)


2m T Error differences from the CTL

T925 mean_abs[CY37R1_CTESSEL(ficd)+36-AN(ficd)]-mean_abs[CY37R1(fhrrd)+36-AN(fhrrd)]



2m Rh Error differences from the CTL

RH mean_abs[CY37R1_CTESSEL(ficd)+36-AN(ficd)]-mean_abs[CY37R1(fhrrd)+36-AN(fhrrd)]



Having better LE/H heat flux from the surface does not always lead to a better atmospheric prediction → interaction with other processes and compensating errors?

Modelling stomatal conductance (empirical vs mechanistic approaches):

$$E = \frac{\beta}{r_c + r_a} (q_a - q_{sat})$$

The Jarvis (statistical) approach
CHTESSEL in IFS (operational)

$$r_c = \frac{r_{S,\min}}{LAI} f_1(R_s) f_2(\bar{\theta}) f_3(D_a)$$

The mechanistic approach
CTESSEL in IFS


$$r_c = f(r_{cc})$$

$$r_{cc} = \frac{\alpha}{A_n} (C_s - C_i)$$

Copernicus
atmospheric CO₂
forecast/analysis

Aspects	Jarvis model	CTESSEL model
Simplicity/robustness	Yes	No
Coupling with carbon cycle & ecosystem CO ₂ flux	No	Yes
Feedbacks on vegetation	No	Yes
Use carbon observations	LAI	LAI, SIF, GPP, atmospheric CO ₂ for mass balance

Feedbacks from vegetation: Impact of assimilating LAI on 2m temperature





NRT_LAI_ALB – FCLIM:

November 2010

**Severe drought in the Horn
of Africa**

Feedbacks from vegetation: Impact of assimilating LAI on albedo









Reduction of cold/moist bias in 3-day FC over northern Europe in March 2015

Impact of dynamic vegetation on monthly forecast in semi-arid regions

Improved skill of monthly forecast 2m-T with soil moisture and dynamic phenology compared to fc with climatologies

Hot-spots of NEE and GPP variability
NEE (DGVM)




Koster and Walker (2015)

Jung et al. JGR 2011

Using carbon observations to improve carbon and NWP: Fluorescence as a proxy for GPP


During photosynthesis a plant absorbs Photosynthetically Active Radiation (PAR) through its chlorophyll:

- % for ecosystem GPP
- % lost as heat
- % re-emitted as chlorophyll fluorescence (SIF)



How light energy falling on a leaf is partitioned. About 78% of the incident radiation is absorbed, while the rest is either transmitted or reflected at the leaf's surface. About 20% is dissipated through heat and only 2% emitted as fluorescence, as a by-product of photosynthetic reactions occurring within the leaf itself.


A simpler approach with a statistical model



Relationship between GPP
and SIF is ~ linear

$R^2(\text{SIF, GPP})$
=0.8

$$y = -0.88 + 3.55x; r^2 = 0.92$$
$$y = 0.35 + 3.71x; r^2 = 0.79$$
$$y = -0.17 + 3.48x; r^2 = 0.87$$



Guanter et al. (2014)

- $\text{GPP} = a + b \times \text{SIF}$
- a & b coefficients function of PFTs
-


Mac Bean et al. in prep.

Transpiration of water vapour from plants is correlated with CO₂ uptake (GPP)




Figure 36-3 Biological Science, 2/e
© 2005 Pearson Prentice Hall, Inc.

$$ET = \frac{GPP}{WUE}$$



Improving GPP and WUE in models should lead to a better ET







Feedbacks of carbon cycle to NWP:

- Thermal infrared radiative transfer in model and data assimilation**

Radiative forcing of greenhouse gases




Myhre, Shindell et al. (2015) IPCC report AR5, Chapter 8

Shortwave: atmosphere is mostly transparent
Longwave: atmosphere is mostly opaque


Using variable CO₂ for the assimilation of the thermal IR

Reduction of bias correction in varBC: IASI channel ~ 700 hPa

(a) VarBC correction with fixed CO₂



(b) VarBC correction with variable CO₂ from MACC





Atmospheric CH₄ in the ECMWF model (IFS)

CH₄ synoptic variability: 25 to 29th of March 2010

Average total column CH₄ [ppb]

Mid-tropospheric CH₄ [ppb] at 400 hPa



Chemical production of water vapour : CH₄ oxidation



Parameterization in IFS:

$$\Delta[\text{H}_2\text{O}] = 2k_1[\text{CH}_4]$$

$$\Delta[\text{H}_2\text{O}] = k_1(6.8 - [\text{H}_2\text{O}])$$

Simmons, Randel et al. 1998,
Brasseur and Solomon 1984
Monge-Sanz et al. 2013

- Change of CH₄ associated with transport and global CH₄ increase no considered.
- Assumption breaks in polar regions (removal of H₂O by condensation).



Randel et al. 1998

Summary

- **Carbon cycle is at the heart of climate change (long time scales > 1year)**

Climatologies of atmospheric composition in NWP


- **Processes on shorter time-scales relevant for NWP (1-day to 1-year):**

Dynamic vegetation model to link water, energy and carbon cycles.




Explore impact on skill for long (**monthly, seasonal**) and **high resolution** forecasts?

- **Copernicus Atmosphere Monitoring Service future work on carbon cycle could benefit NWP:**
 - Explore use of chlorophyll fluorescence retrievals from satellites to evaluate/constrain photosynthesis in the model (impact on carbon, water and energy fluxes).
 - Score carbon, water and energy fluxes using eddy covariance observations in near-real time

20100101



[ppm]



Thank you



Implemented by ECMWF

S. Massart